

ASYMPTOTICS FOR RANDOMLY REINFORCED URNS WITH RANDOM BARRIERS

PATRIZIA BERTI, IRENE CRIMALDI, LUCA PRATELLI, AND PIETRO RIGO

ABSTRACT. An urn contains black and red balls. Let Z_n be the proportion of black balls at time n and $0 \leq L < U \leq 1$ random barriers. At each time n , a ball b_n is drawn. If b_n is black and $Z_{n-1} < U$, then b_n is replaced together with a random number B_n of black balls. If b_n is red and $Z_{n-1} > L$, then b_n is replaced together with a random number R_n of red balls. Otherwise, no additional balls are added, and b_n alone is replaced. In this paper, we assume $R_n = B_n$. Then, under mild conditions, it is shown that $Z_n \xrightarrow{a.s.} Z$ for some random variable Z , and

$$D_n := \sqrt{n}(Z_n - Z) \longrightarrow \mathcal{N}(0, \sigma^2) \quad \text{conditionally a.s.}$$

where σ^2 is a certain random variance. Almost sure conditional convergence means that

$$P(D_n \in \cdot \mid \mathcal{G}_n) \xrightarrow{weakly} \mathcal{N}(0, \sigma^2) \quad \text{a.s.}$$

where $P(D_n \in \cdot \mid \mathcal{G}_n)$ is a regular version of the conditional distribution of D_n given the past $\mathcal{G}_n$. Thus, in particular, one obtains $D_n \longrightarrow \mathcal{N}(0, \sigma^2)$ stably. It is also shown that $L < Z < U$ a.s. and Z has non-atomic distribution.

1. INTRODUCTION

In recent times, there is a growing interest on randomly reinforced urns. A meaningful version of the latter, introduced in [2] and supported by real applications, is the following.

1.1. Framework. An urn contains $b > 0$ black balls and $r > 0$ red balls. At each time, a ball is drawn and then replaced, possibly together with a random number of balls of the same color. Precisely, for each $n \geq 1$, let b_n denote the ball drawn at time n and Z_n the proportion of black balls in the urn at time n . Then,

- If b_n is black and $Z_{n-1} < U$, where U is a random barrier, b_n is replaced together with a random number $B_n \geq 0$ of black balls;
- If b_n is red and $Z_{n-1} > L$, where $L < U$ is another random barrier, b_n is replaced together with a random number $R_n \geq 0$ of red balls;
- Otherwise, b_n is replaced without additional balls, so that the composition of the urn does not change.

Date: August 27, 2015.

2010 Mathematics Subject Classification. 60B10, 60F05, 60G57, 62F15.

Key words and phrases. Bayesian nonparametrics – Central limit theorem – Clinical trial – Random probability measure – Stable convergence – Urn model.

To model such urns, we fix a probability space $(\Omega, \mathcal{A}, P)$ supporting the random variables $(L, U, X_n, B_n, R_n : n \geq 1)$ such that

$$0 \leq L < U \leq 1; \quad X_n \in \{0, 1\}; \quad 0 \leq B_n, R_n \leq c \quad \text{for some constant } c.$$

We let

$$\begin{aligned} \mathcal{G}_0 &= \sigma(L, U), \quad \mathcal{G}_n = \sigma(L, U, X_1, B_1, R_1, \dots, X_n, B_n, R_n), \quad Z_0 = b/(b+r), \\ Z_n &= \frac{b + \sum_{i=1}^n X_i B_i I_{\{Z_{i-1} < U\}}}{b + r + \sum_{i=1}^n [X_i B_i I_{\{Z_{i-1} < U\}} + (1 - X_i) R_i I_{\{Z_{i-1} > L\}}]}. \end{aligned}$$

Further, we assume

$$\begin{aligned} E(X_{n+1} \mid \mathcal{G}_n) &= Z_n \text{ a.s.} \quad \text{and} \\ (B_n, R_n) &\text{ independent of } \sigma(\mathcal{G}_{n-1}, X_n). \end{aligned}$$

Clearly, X_n should be regarded as the indicator of the event {black ball at time n } and Z_n as the proportion of black balls in the urn at time n .

1.2. State of the art. Though the literature on randomly reinforced urns is quite huge, random barriers are not so popular. In other terms, the case $L = 0$ and $U = 1$ is widely investigated (see e.g. [1], [3]-[5], [7], [10]-[14] and references therein) but

$$P(\{L > 0\} \cup \{U < 1\}) > 0$$

is almost neglected. To our knowledge, the only explicit reference is [2]. In such a paper, the barriers L and U are not random (i.e., they are constant) and (B_n) and (R_n) are independent sequences of i.i.d. random variables. Then, it is shown that $Z_n \xrightarrow{a.s.} L$ if $E(B_1) < E(R_1)$ and $Z_n \xrightarrow{a.s.} U$ if $E(B_1) > E(R_1)$. Moreover, in case $E(B_1) = E(R_1)$, a method for estimating the unknown mean is proposed. The method is based on $Z_n \xrightarrow{a.s.} Z$, where the random variable Z has non-atomic distribution, if $E(B_1) = E(R_1)$. However, convergence of Z_n when $E(B_1) = E(R_1)$ is stated without a proof. In addition, the limiting distribution of $Z_n - Z$ is not investigated at all.

1.3. Results. In a sense, this paper deals with the opposite case with respect to [2]. Indeed, while (B_n) and (R_n) are independent sequences in [2], throughout this paper it is assumed that

$$R_n = B_n \quad \text{for each } n \geq 1.$$

Under this assumption, the following two results are proved.

Theorem 1. *In the framework of Subsection 1.1, suppose*

$$R_n = B_n \quad \text{and} \quad \liminf_n E(B_n) > 0.$$

Then,

$$Z_n \xrightarrow{a.s.} Z$$

for some random variable Z such that $L \leq Z \leq U$ and $0 < Z < 1$ a.s.

Theorem 2. *In the framework of Subsection 1.1, suppose*

$$R_n = B_n, \quad m := \lim_n E(B_n) > 0 \quad \text{and} \quad q := \lim_n E(B_n^2).$$

Define

$$D_n = \sqrt{n}(Z_n - Z) \quad \text{and} \quad \sigma^2 = qZ(1-Z)/m^2,$$

where Z is the a.s. limit of Z_n . Then,

$$D_n \longrightarrow \mathcal{N}(0, \sigma^2) \quad \text{conditionally a.s. with respect to } (\mathcal{G}_n).$$

Moreover, Z has a non-atomic distribution and $L < Z < U$ a.s.

In Theorem 2, $\mathcal{N}(a, b)$ denotes the Gaussian law with mean a and variance $b \geq 0$, where $\mathcal{N}(a, 0) = \delta_a$. Almost sure conditional convergence is a strong form of stable convergence, introduced in [8]-[9] and involved in [1], [5], [13], [14]. The general definition is discussed in Section 2. In the present case, it means that

$$P(D_n \in \cdot \mid \mathcal{G}_n)(\omega) \xrightarrow{\text{weakly}} \mathcal{N}(0, \sigma^2(\omega)) \quad \text{for almost all } \omega \in \Omega$$

where $P(D_n \in \cdot \mid \mathcal{G}_n)$ is a regular version of the conditional distribution of D_n given $\mathcal{G}_n$. Thus, in particular, Theorem 2 yields

$$D_n \longrightarrow \mathcal{N}(0, \sigma^2) \quad \text{stably;}$$

see Lemma 3.

Theorems 1-2 establish the asymptotics for randomly reinforced urns with random barriers when $R_n = B_n$. The case $R_n \neq B_n$, as well as some other possible developments, are discussed in Section 4. A last note is that Theorem 2 agrees with the result obtained when random barriers are not taken into account. Indeed, if $L = 0$ and $U = 1$, Theorem 2 follows from [5, Corollary 3].

2. ALMOST SURE CONDITIONAL CONVERGENCE

Almost sure conditional convergence, introduced in [8]-[9], may be regarded as a strong form of stable convergence. We now make it precise.

Let $(\Omega, \mathcal{A}, P)$ be a probability space and S a metric space. A *kernel* on S (or a *random probability measure* on S) is a measurable collection $N = \{N(\omega) : \omega \in \Omega\}$ of probability measures on the Borel σ -field on S . Measurability means that

$$N(\cdot)(f) = \int f(x) N(\cdot)(dx)$$

is a real random variable for each bounded Borel map $f : S \rightarrow \mathbb{R}$. To denote such random variable, in the sequel, we will often write $N(f)$ instead of $N(\cdot)(f)$.

For each $n \geq 1$, fix a sub- σ -field $\mathcal{F}_n \subset \mathcal{A}$. Also, let (Y_n) be a sequence of S -valued random variables and N a kernel on S . Say that Y_n converges to N , *conditionally a.s. with respect to* $(\mathcal{F}_n)$, if

$$E\{f(Y_n) \mid \mathcal{F}_n\} \xrightarrow{\text{a.s.}} N(f) \quad \text{for each } f \in C_b(S). \quad (1)$$

If S is Polish, condition (1) has a quite transparent meaning. Suppose in fact S is Polish and fix a regular version $P(Y_n \in \cdot \mid \mathcal{F}_n)$ of the conditional distribution of Y_n given $\mathcal{F}_n$. Then, condition (1) is equivalent to

$$P(Y_n \in \cdot \mid \mathcal{F}_n)(\omega) \xrightarrow{\text{weakly}} N(\omega) \quad \text{for almost all } \omega \in \Omega.$$

So far, $(\mathcal{F}_n)$ is an arbitrary sequence of sub- σ -fields. Suppose now that $(\mathcal{F}_n)$ is a filtration, in the sense that $\mathcal{F}_n \subset \mathcal{F}_{n+1} \subset \mathcal{A}$ for each n . Then, under a mild measurability condition, almost sure conditional convergence implies stable

convergence. This is noted in [8, Section 5] but we give a proof to make the paper self-contained. Let

$$\mathcal{F}_\infty = \sigma(\cup_n \mathcal{F}_n).$$

Lemma 3. *Suppose $(\mathcal{F}_n)$ is a filtration such that*

$$N(f) \text{ and } Y_n \text{ are } \mathcal{F}_\infty\text{-measurable for all } f \in C_b(S) \text{ and } n \geq 1.$$

If $Y_n \rightarrow N$ conditionally a.s. with respect to $(\mathcal{F}_n)$, then $Y_n \rightarrow N$ stably, that is

$$E\{N(f) \mid H\} = \lim_n E\{f(Y_n) \mid H\}$$

whenever $f \in C_b(S)$, $H \in \mathcal{A}$ and $P(H) > 0$.

Proof. Let $f \in C_b(S)$ and $H \in \mathcal{A}$. If $H \in \cup_n \mathcal{F}_n$, then $H \in \mathcal{F}_n$ for each sufficiently large n , so that

$$E\{N(f) I_H\} = \lim_n E\left(E\{f(Y_n) \mid \mathcal{F}_n\} I_H\right) = \lim_n E\{f(Y_n) I_H\}.$$

Since $\cup_n \mathcal{F}_n$ is a field, by standard arguments one obtains

$$E\{N(f) V\} = \lim_n E\{f(Y_n) V\} \quad \text{whenever } V \text{ is bounded and } \mathcal{F}_\infty\text{-measurable.}$$

Hence, for arbitrary $H \in \mathcal{A}$, the measurability condition implies

$$E\{N(f) I_H\} = E\left(N(f) E(I_H \mid \mathcal{F}_\infty)\right) = \lim_n E\left(f(Y_n) E(I_H \mid \mathcal{F}_\infty)\right) = \lim_n E\{f(Y_n) I_H\}.$$

□

Note that the measurability condition of Lemma 3 is trivially true if $\mathcal{F}_\infty = \mathcal{A}$.

We refer to [8]-[9] for more on almost sure conditional convergence. Here, for easy of reference, we report three useful facts. The first and the second are already known (see [5, Proposition 1 and Lemma 2] and [9, Theorem 2.2]) while the third is a quick consequence of condition (1). In each of these facts, $(\mathcal{F}_n)$ is a filtration.

Lemma 4. *Suppose the Y_n are real random variables such that $Y_n \xrightarrow{a.s.} Y$. Then,*

$$\sqrt{n}(Y_n - Y) \longrightarrow \mathcal{N}(0, U), \quad \text{conditionally a.s. with respect to } (\mathcal{F}_n),$$

where U is a real random variable, provided

- (i) (Y_n) is a uniformly integrable martingale with respect to $(\mathcal{F}_n)$;
- (ii) $E\{\sup_n \sqrt{n}|Y_n - Y_{n-1}|\} < \infty$;
- (iii) $n \sum_{k \geq n} (Y_k - Y_{k-1})^2 \xrightarrow{a.s.} U$.

Lemma 5. *Suppose the Y_n are real random variables. If (Y_n) is adapted to $(\mathcal{F}_n)$, $\sum_n n^{-2} E(Y_n^2) < \infty$ and $E(Y_{n+1} \mid \mathcal{F}_n) \xrightarrow{a.s.} U$, for some real random variable U , then*

$$n \sum_{k \geq n} \frac{Y_k}{k^2} \xrightarrow{a.s.} U \quad \text{and} \quad \frac{1}{n} \sum_{k=1}^n Y_k \xrightarrow{a.s.} U.$$

Lemma 6. *Suppose $Y_n \rightarrow N$ conditionally a.s. with respect to $(\mathcal{F}_n)$. Define $Q(A) = E\{I_A V\}$ for $A \in \mathcal{A}$, where $V \geq 0$, $E(V) = 1$ and V is $\mathcal{F}_\infty$ -measurable. Then $Y_n \rightarrow N$, conditionally a.s. with respect to $(\mathcal{F}_n)$, under Q as well.*

Proof. Suppose first $\sup V < \infty$ and define $K_n = V - E(V \mid \mathcal{F}_n)$. Given $f \in C_b(S)$,

$$E_Q\{f(Y_n) \mid \mathcal{F}_n\} = \frac{E\{V f(Y_n) \mid \mathcal{F}_n\}}{E(V \mid \mathcal{F}_n)} = E\{f(Y_n) \mid \mathcal{F}_n\} + \frac{E\{K_n f(Y_n) \mid \mathcal{F}_n\}}{E(V \mid \mathcal{F}_n)}, \quad Q\text{-a.s.},$$

where E_Q denotes expectation under Q . Since $\sigma(V) \subset \mathcal{F}_\infty$ and $|K_n| \leq \sup V$ a.s., the martingale convergence theorem (in the version of [6]) implies

$$E(V \mid \mathcal{F}_n) \xrightarrow{a.s.} E(V \mid \mathcal{F}_\infty) = V \quad \text{and} \\ \left| E\{K_n f(Y_n) \mid \mathcal{F}_n\} \right| \leq \sup |f| E\{|K_n| \mid \mathcal{F}_n\} \xrightarrow{a.s.} 0.$$

Since $Q(V > 0) = 1$, one obtains $E_Q\{f(Y_n) \mid \mathcal{F}_n\} \rightarrow N(f)$, Q -a.s. This concludes the proof for bounded V . If V is not bounded, it suffices to reply V with $V I_{\{V \leq v\}}/E(V I_{\{V \leq v\}})$ and to take the limit as $v \rightarrow \infty$. $\square$

3. PROOFS

In the sequel, for any events $A_n \in \mathcal{A}$ and $B \in \mathcal{A}$, we say that A_n is *eventually true on* B (or, more briefly, A_n *eventually on* B) whenever

$$P\{\omega \in B : \omega \notin A_n \text{ for infinitely many } n\} = 0.$$

Assume the conditions of Subsection 1.1 and $R_n = B_n$. Let

$$S_n = b + r + \sum_{i=1}^n [X_i B_i I_{\{Z_{i-1} < U\}} + (1 - X_i) B_i I_{\{Z_{i-1} > L\}}]$$

denote the denominator of Z_n , namely, the number of balls in the urn at time n . Also, the filtration $(\mathcal{G}_n)$ is abbreviated by $\mathcal{G}$.

After some (tedious but easy) algebra, one obtains

$$Z_{n+1} - Z_n = Z_n H_n + \Delta_{n+1},$$

where

$$H_n = \frac{B_{n+1}}{S_n + B_{n+1}} (1 - Z_n) (I_{\{Z_n < U\}} - I_{\{Z_n > L\}}), \\ \Delta_{n+1} = \frac{B_{n+1}}{S_n + B_{n+1}} (X_{n+1} - Z_n) ((1 - Z_n) I_{\{Z_n < U\}} + Z_n I_{\{Z_n > L\}}).$$

This writing of $Z_{n+1} - Z_n$ is fundamental for our purposes.

3.1. Proof of Theorem 1. In this subsection, it is assumed that

$$\liminf_n E(B_n) > 0.$$

Since

$$E(X_{n+1} \mid \mathcal{G}_n, B_{n+1}) = E(X_{n+1} \mid \mathcal{G}_n) = Z_n \quad \text{a.s.},$$

then

$$E(\Delta_{n+1} \mid \mathcal{G}_n) = 0 \quad \text{a.s.}$$

This fact has two useful consequences. First,

$$M_n = \sum_{i=1}^n \Delta_i$$

is a $\mathcal{G}$ -martingale. Second, (Z_n) is a $\mathcal{G}$ -sub-martingale in case $U = 1$. In fact, $U = 1$ implies $H_n \geq 0$, so that

$$E\{Z_{n+1} - Z_n \mid \mathcal{G}_n\} = E\{Z_n H_n + \Delta_{n+1} \mid \mathcal{G}_n\} = Z_n E(H_n \mid \mathcal{G}_n) \geq 0 \quad \text{a.s.}$$

Similarly, if $L = 0$ then (Z_n) is a $\mathcal{G}$ -super-martingale. Therefore, it is not hard to see that Z_n converges a.s. on the set $\{L = 0\} \cup \{U = 1\}$.

We next state two lemmas.

Lemma 7. *Let $Z_* = \liminf_n Z_n$ and $Z^* = \limsup_n Z_n$. Each of the following statements implies the subsequent:*

- (a) $0 < L < U < 1$ a.s.;
- (b) $0 < Z_* \leq Z^* < 1$ a.s.;
- (c) $\liminf_n (S_n/n) > 0$ a.s.;
- (d) M_n converges a.s.

Proof. “(a) $\implies$ (b)”. Let $H = \{Z_* = 0, L > 0\}$. On the set H , one obtains

$$\sup_n S_n = \infty, \quad \lim_n (Z_{n+1} - Z_n) = 0, \quad Z_n > L \text{ for infinitely many } n.$$

Define $\tau_0 = 0$ and

$$\tau_n = \inf\{k : k > \tau_{n-1}, \quad Z_{k-1} > L, \quad Z_k \leq L\}.$$

Then, $\tau_n < \infty$ for all n on H . Observe now that $Z_j \geq Z_{j-1}$ whenever $Z_{j-1} \leq L$. Hence, $Z_* = \liminf_n Z_n = \liminf_n Z_{\tau_n}$ on H , which implies the contradiction

$$Z_* \geq \liminf_n Z_{\tau_{n-1}} + \liminf_n (Z_{\tau_n} - Z_{\tau_{n-1}}) = \liminf_n Z_{\tau_{n-1}} \geq L > 0 \quad \text{a.s. on } H.$$

Thus, under (a), one obtains $P(Z_* = 0) = P(H) = 0$. Similarly, $P(Z^* = 1) = 0$.

“(b) $\implies$ (c)”. Define

$$K_n = \sum_{i=1}^n \frac{I_{\{Z_{i-1} < U\}} [X_i B_i - Z_{i-1} E(B_i)] + I_{\{Z_{i-1} > L\}} [(1 - X_i) B_i - (1 - Z_{i-1}) E(B_i)]}{i}.$$

Since K_n is a $\mathcal{G}$ -martingale and $\sup_n E(K_n^2) < \infty$, then K_n converges a.s. Thus, Kronecker lemma implies $(1/n) \sum_{i=1}^n i K_i \xrightarrow{\text{a.s.}} 0$, so that

$$\liminf_n \frac{S_n}{n} = \liminf_n \frac{1}{n} \sum_{i=1}^n \left(I_{\{Z_{i-1} < U\}} Z_{i-1} E(B_i) + I_{\{Z_{i-1} > L\}} (1 - Z_{i-1}) E(B_i) \right) \text{ a.s.}$$

Since $I_{\{Z_{i-1} < U\}} + I_{\{Z_{i-1} > L\}} \geq 1$, one finally obtains

$$\liminf_n \frac{S_n}{n} \geq \{Z_* \wedge (1 - Z^*)\} \liminf_n E(B_n) > 0 \quad \text{a.s.}$$

“(c) $\implies$ (d)”. Since $0 \leq B_{n+1} \leq c$,

$$E\{(M_{n+1} - M_n)^2 \mid \mathcal{G}_n\} = E(\Delta_{n+1}^2 \mid \mathcal{G}_n) \leq E\left(\frac{B_{n+1}^2}{S_n^2} \mid \mathcal{G}_n\right) = \frac{E(B_{n+1}^2)}{S_n^2} \leq \frac{c^2}{n^2} \frac{1}{(S_n/n)^2} \quad \text{a.s.}$$

Thus, $\sum_n E\{(M_{n+1} - M_n)^2 \mid \mathcal{G}_n\} < \infty$ a.s. by condition (c). It follows that the $\mathcal{G}$ -martingale M_n converges a.s. $\square$

Lemma 8. *If $\liminf_n (S_n/n) > 0$ a.s., then $P(D) = 0$ where*

$$D = \{Z_a \leq L \text{ for infinitely many } a \text{ and } Z_b \geq U \text{ for infinitely many } b\}.$$

Proof. On D , there is a sequence (a_n, b_n) such that $a_1 < b_1 < a_2 < b_2 < \dots$ and

$$Z_{a_n} \leq L, \quad Z_{b_n} \geq U, \quad L < Z_k < U \text{ for each } a_n < k < b_n.$$

Since $H_k = 0$ if $a_n < k < b_n$, then

$$U - L \leq Z_{b_n} - Z_{a_n} = \sum_{k=a_n}^{b_n-1} (Z_{k+1} - Z_k) = \sum_{k=a_n}^{b_n-1} (Z_k H_k + \Delta_{k+1}) = Z_{a_n} H_{a_n} + M_{b_n} - M_{a_n}.$$

Since $\liminf_n (S_n/n) > 0$ a.s., then $\sup_n S_n = \infty$ a.s., which implies $H_n \xrightarrow{a.s.} 0$. Also, by Lemma 7, M_n converges a.s. Hence, taking the limit as $n \rightarrow \infty$, one obtains $U - L \leq 0$ a.s. on D . Therefore, $P(D) = 0$. $\square$

We are now ready to prove a.s. convergence of Z_n .

Since Z_n converges a.s. on the set $\{L = 0\} \cup \{U = 1\}$, it can be assumed $P(0 < L < U < 1) > 0$. In turn, up to replacing P with $P(\cdot \mid 0 < L < U < 1)$, it can be assumed $0 < L < U < 1$ a.s. Then, Lemmas 7-8 imply $P(D^c) = 1$ and a.s. convergence of M_n . Write

$$Z_n - Z_0 = \sum_{i=0}^{n-1} (Z_{i+1} - Z_i) = \sum_{i=0}^{n-1} Z_i H_i + M_n = K_n + M_n$$

where $K_n = \sum_{i=0}^{n-1} Z_i H_i$. On the set D^c , one has either $Z_i H_i \geq 0$ eventually or $Z_i H_i \leq 0$ eventually. Hence, on D^c , the sequence K_n converges if and only if it is bounded. But K_n is a.s. bounded, since $|K_n| \leq 1 + \sup_k |M_k|$ and M_n converges a.s. Thus, Z_n converges a.s. on D^c . This proves a.s. convergence of Z_n for $P(D^c) = 1$.

Let Z denote the a.s. limit of Z_n . Since $Z_n \xrightarrow{a.s.} 1$ on the set $\{Z < L\}$ and $Z_n \xrightarrow{a.s.} 0$ on the set $\{Z > U\}$, then $L \leq Z \leq U$ a.s.

It remains to see that

$$P(Z = 0) = P(Z = 1) = 0.$$

We just prove $P(Z = 1) = 0$. The proof of $P(Z = 0) = 0$ is quite analogous.

Since $Z \leq U \leq 1$ a.s., then $P(Z = 1) \leq P(U = 1)$. Thus, it can be assumed $P(U = 1) > 0$. In turn, up to replacing P with $P(\cdot \mid U = 1)$, it can be assumed $U = 1$ everywhere. Then, Z_n is a $\mathcal{G}$ -sub-martingale, so that

$$Y_n = Z_n / (1 - Z_n)$$

is still a $\mathcal{G}$ -sub-martingale. Let $H = \{\sum_n E\{Y_{n+1} - Y_n \mid \mathcal{G}_n\} < \infty\}$. Since Y_n is a positive $\mathcal{G}$ -sub-martingale, Y_n converges a.s. (to a real random variable) on the set H . Thus, to get $P(Z = 1) = 0$, it suffices to show that

$$\sum_n E\{Y_{n+1} - Y_n \mid \mathcal{G}_n\} < \infty \quad \text{a.s. on the set } \{Z = 1\}. \quad (2)$$

To prove (2), let

$$J_n = b + \sum_{i=1}^n X_i B_i \quad \text{and} \quad L_n = S_n - J_n = r + \sum_{i=1}^n (1 - X_i) B_i I_{\{Z_{i-1} > L\}}$$

be the numbers of black balls and red balls, respectively, in the urn at time n (recall that $U = 1$, so that $Z_{i-1} < U$ is automatically true). On noting that $Y_n = J_n/L_n$, one obtains

$$\begin{aligned}
E\{Y_{n+1} - Y_n \mid \mathcal{G}_n\} &= -Y_n + E\left\{\frac{J_n + B_{n+1}}{L_n} X_{n+1} + \frac{J_n}{L_n + I_{\{Z_n > L\}} B_{n+1}} (1 - X_{n+1}) \mid \mathcal{G}_n\right\} \\
&= -Y_n + Z_n E\left\{\frac{J_n + B_{n+1}}{L_n} \mid \mathcal{G}_n\right\} + (1 - Z_n) E\left\{\frac{J_n}{L_n + I_{\{Z_n > L\}} B_{n+1}} \mid \mathcal{G}_n\right\} \\
&= -Y_n (1 - Z_n) + \frac{Z_n E(B_{n+1})}{L_n} + Y_n (1 - Z_n) E\left\{\frac{L_n}{L_n + I_{\{Z_n > L\}} B_{n+1}} \mid \mathcal{G}_n\right\} \\
&= \frac{Z_n E(B_{n+1})}{L_n} - Z_n I_{\{Z_n > L\}} E\left\{\frac{B_{n+1}}{L_n + I_{\{Z_n > L\}} B_{n+1}} \mid \mathcal{G}_n\right\} \\
&\leq \frac{Z_n E(B_{n+1})}{L_n} - Z_n I_{\{Z_n > L\}} E\left\{\frac{B_{n+1}}{L_n + c} \mid \mathcal{G}_n\right\} \\
&= \frac{Z_n E(B_{n+1})}{L_n} - Z_n I_{\{Z_n > L\}} \frac{E(B_{n+1})}{L_n + c} \quad \text{a.s.}
\end{aligned}$$

Since $Z_n \xrightarrow{\text{a.s.}} Z$, then $Z_n > L$ eventually on the set $\{Z = 1\}$. Hence,

$$E\{Y_{n+1} - Y_n \mid \mathcal{G}_n\} \leq Z_n E(B_{n+1}) \left(\frac{1}{L_n} - \frac{1}{L_n + c}\right) \leq \frac{c^2}{L_n^2} \quad \text{eventually on } \{Z = 1\}.$$

Next, given $k \in (1, 2)$, it is not hard to see that

$$E\left\{\frac{J_{n+1}}{L_{n+1}^k} - \frac{J_n}{L_n^k} \mid \mathcal{G}_n\right\} \leq 0 \quad \text{eventually on } \{Z = 1\}.$$

We omit the calculations for they exactly agree with those for proving [12, Lemma A.1(ii)]. Thus, the sequence J_n/L_n^k converges a.s. on $\{Z = 1\}$. Furthermore, independence of the B_n yields

$$\begin{aligned}
\liminf_n \frac{J_n}{n} &= \liminf_n \frac{\sum_{i=1}^n B_i}{n} \frac{S_n}{\sum_{i=1}^n B_i} Z_n = \liminf_n \frac{\sum_{i=1}^n B_i}{n} \\
&= \liminf_n \frac{\sum_{i=1}^n E(B_i)}{n} \geq \liminf_n E(B_n) > 0 \quad \text{a.s. on } \{Z = 1\}.
\end{aligned}$$

Given any $\gamma < 1$, it follows that

$$\frac{n^\gamma}{L_n^k} = \frac{n^\gamma}{n} \frac{n}{J_n} \frac{J_n}{L_n^k} \xrightarrow{\text{a.s.}} 0 \quad \text{a.s. on } \{Z = 1\}.$$

Thus,

$$L_n > n^{\gamma/k} \quad \text{eventually on } \{Z = 1\}.$$

Since $k < 2$, one can take $\gamma < 1$ such that $\gamma/k > 1/2$. Therefore, condition (2) holds, and this concludes the proof of Theorem 1.

3.2. Proof of Theorem 2. In this subsection, it is assumed that

$$m := \lim_n E(B_n) > 0 \quad \text{and} \quad q := \lim_n E(B_n^2).$$

By Theorem 1, $Z_n \xrightarrow{\text{a.s.}} Z$ for some random variable Z such that $L \leq Z \leq U$ and $0 < Z < 1$ a.s.

On noting that

$$0 < Z_* = Z = Z^* < 1 \quad \text{a.s.},$$

the same argument used after Lemma 8 yields $\sum_n Z_n |H_n| < \infty$ a.s. Since $Z > 0$ a.s., it follows that

$$\sum_n |H_n| < \infty \quad \text{a.s.} \quad (3)$$

Define

$$T_n = \prod_{i=1}^{n-1} (1 + H_i) \quad \text{and} \quad W_n = \frac{Z_n}{T_n}.$$

Condition (3) implies $T_n \xrightarrow{a.s.} T$, for some real random variable $T > 0$, so that

$$W_n \xrightarrow{a.s.} Z/T := W.$$

Our next goal is to show that $\sqrt{n}(W_n - W)$ converges conditionally a.s. To this end, we first fix the asymptotic behavior of S_n .

Lemma 9. $S_n/n \xrightarrow{a.s.} m$.

Proof. Let $Q_n = \sum_{i=1}^n B_i [X_i I_{\{Z_{i-1} \geq U\}} + (1 - X_i) I_{\{Z_{i-1} \leq L\}}]$. Since $Z_i \xrightarrow{a.s.} Z$, then $1 - Z_i > (1 - Z)/2$ eventually. Moreover, $I_{\{Z_{i-1} \geq U\}} + I_{\{Z_{i-1} \leq L\}} = |I_{\{Z_{i-1} < U\}} - I_{\{Z_{i-1} > L\}}|$. Therefore,

$$\begin{aligned} B_i [X_i I_{\{Z_{i-1} \geq U\}} + (1 - X_i) I_{\{Z_{i-1} \leq L\}}] &\leq B_i |I_{\{Z_{i-1} < U\}} - I_{\{Z_{i-1} > L\}}| \\ &= |H_{i-1}| \frac{S_{i-1} + B_i}{1 - Z_{i-1}} \leq \frac{2|H_{i-1}|}{1 - Z} (S_{i-1} + B_i) \quad \text{eventually.} \end{aligned}$$

By condition (3) and Kronecker lemma,

$$\frac{Q_n}{S_n} \leq \frac{2}{1 - Z} \frac{1}{S_n} \sum_{i=1}^n |H_{i-1}| S_{i-1} + \frac{2c}{1 - Z} \frac{1}{S_n} \sum_{i=1}^n |H_{i-1}| \xrightarrow{a.s.} 0.$$

Hence,

$$\frac{Q_n}{n} = \frac{Q_n}{S_n} \frac{S_n}{n} \leq \frac{Q_n}{S_n} \frac{r + b + nc}{n} \xrightarrow{a.s.} 0.$$

On noting that $(1/n) \sum_{i=1}^n B_i \xrightarrow{a.s.} m$, one finally obtains

$$\frac{S_n}{n} = \frac{r + b}{n} - \frac{Q_n}{n} + \frac{\sum_{i=1}^n B_i}{n} \xrightarrow{a.s.} m.$$

□

In view of the next lemma, we recall that

$$\sigma^2 = q Z (1 - Z) / m^2.$$

Lemma 10.

$$\sqrt{n}(W_n - W) \longrightarrow \mathcal{N}(0, \sigma^2/T^2) \quad \text{conditionally a.s. with respect to } \mathcal{G}.$$

Proof. First note that W_n can be written as

$$W_n = Z_1 + \sum_{i=1}^{n-1} \frac{\Delta_{i+1}}{T_{i+1}}.$$

Thus, W_n is a $\mathcal{G}$ -martingale and the obvious strategy would be applying Lemma 4 to $Y_n = W_n$. However, conditions (i)-(ii)-(iii) are not easy to check with $Y_n = W_n$. Accordingly, we adopt an approximation procedure.

Given $\epsilon > 0$, define

$$W_n^{(\epsilon)} = Z_1 + \sum_{i=1}^{n-1} \frac{\Delta_{i+1} I_{A_i}}{\epsilon \vee T_{i+1}} \quad \text{where } A_i = \{2 S_i > i m\}.$$

For fixed $\epsilon > 0$, $W_n^{(\epsilon)}$ is still a $\mathcal{G}$ -martingale and

$$\sup_n E\{(W_n^{(\epsilon)})^2\} \leq 1 + \frac{1}{\epsilon^2} \sum_{i=1}^{\infty} E\{\Delta_{i+1}^2 I_{A_i}\} \leq 1 + \frac{c^2}{\epsilon^2} \sum_{i=1}^{\infty} E\left\{\frac{I_{A_i}}{S_i^2}\right\} \leq 1 + \left(\frac{2c}{m\epsilon}\right)^2 \sum_{i=1}^{\infty} \frac{1}{i^2}.$$

Hence, $W_n^{(\epsilon)} \xrightarrow{a.s.} W^{(\epsilon)}$ for some random variable $W^{(\epsilon)}$. Since $S_n/n \xrightarrow{a.s.} m$, the events A_n are eventually true, so that

$$W - W_n = \sum_{i \geq n} \frac{\Delta_{i+1}}{T_{i+1}} = \sum_{i \geq n} \frac{\Delta_{i+1} I_{A_i}}{\epsilon \vee T_{i+1}} = W^{(\epsilon)} - W_n^{(\epsilon)} \quad \text{eventually on } \{T > \epsilon\}.$$

Therefore, it suffices to show that, for fixed $\epsilon > 0$,

$$\sqrt{n} \{W_n^{(\epsilon)} - W^{(\epsilon)}\} \longrightarrow \mathcal{N}(0, \sigma^2/(\epsilon \vee T)^2) \quad \text{conditionally a.s. with respect to } \mathcal{G}.$$

In turn, since $W_n^{(\epsilon)}$ is a uniformly integrable $\mathcal{G}$ -martingale, it suffices to check conditions (ii)-(iii) of Lemma 4 with $Y_n = W_n^{(\epsilon)}$ and $U = \sigma^2/(\epsilon \vee T)^2$. As to (ii),

$$\begin{aligned} E\left\{\left(\sup_n \sqrt{n} |W_n^{(\epsilon)} - W_{n-1}^{(\epsilon)}|\right)^4\right\} &\leq \sum_n n^2 E\{(W_n^{(\epsilon)} - W_{n-1}^{(\epsilon)})^4\} \\ &\leq \frac{1}{\epsilon^4} \sum_n n^2 E\{\Delta_n^4 I_{A_{n-1}}\} \leq \frac{c^4}{\epsilon^4} \sum_n n^2 E\left\{\frac{I_{A_{n-1}}}{S_{n-1}^4}\right\} \leq \left(\frac{2c}{m\epsilon}\right)^4 \sum_n \frac{n^2}{(n-1)^4} < \infty. \end{aligned}$$

We next turn to condition (iii). We have to prove

$$n \sum_{k \geq n} (W_k^{(\epsilon)} - W_{k-1}^{(\epsilon)})^2 = n \sum_{k \geq n} \frac{I_{A_{k-1}} \Delta_k^2}{(\epsilon \vee T_k)^2} \xrightarrow{a.s.} \sigma^2/(\epsilon \vee T)^2.$$

Since $T_k \xrightarrow{a.s.} T$, the above condition reduces to

$$n \sum_{k \geq n} I_{A_{k-1}} \Delta_k^2 \xrightarrow{a.s.} \sigma^2. \quad (4)$$

Since $1 - Z_k > (1 - Z)/2$ eventually and $\sum_k |H_k| < \infty$ a.s., Abel summation formula yields

$$\begin{aligned} & n \sum_{k \geq n} I_{A_{k-1}} \frac{B_k^2}{(S_{k-1} + B_k)^2} \left(I_{\{Z_{k-1} \geq U\}} + I_{\{Z_{k-1} \leq L\}} \right) \\ &= n \sum_{k \geq n} I_{A_{k-1}} \frac{B_k}{S_{k-1} + B_k} \frac{|H_{k-1}|}{1 - Z_{k-1}} \\ &\leq \frac{2cn}{1 - Z} \sum_{k \geq n} I_{A_{k-1}} \frac{|H_{k-1}|}{S_{k-1}} \leq \frac{4cn}{m(1 - Z)} \sum_{k \geq n} \frac{|H_{k-1}|}{k-1} \xrightarrow{a.s.} 0. \end{aligned}$$

Hence, to get (4), it suffices to prove that

$$n \sum_{k \geq n} I_{A_{k-1}} \frac{B_k^2}{(S_{k-1} + B_k)^2} (X_k - Z_{k-1})^2 \xrightarrow{a.s.} \sigma^2.$$

Finally, such condition follows from Lemma 5 if $E(V_{n+1} | \mathcal{G}_n) \xrightarrow{a.s.} \sigma^2$, where

$$V_n = n^2 I_{A_{n-1}} \frac{B_n^2}{(S_{n-1} + B_n)^2} (X_n - Z_{n-1})^2.$$

In fact,

$$\begin{aligned} E(V_{n+1} | \mathcal{G}_n) &= I_{A_n} (n+1)^2 E \left\{ \frac{B_{n+1}^2}{(S_n + B_{n+1})^2} (X_{n+1} - Z_n)^2 | \mathcal{G}_n \right\} \leq (n+1)^2 E \left\{ \frac{B_{n+1}^2}{S_n^2} (X_{n+1} - Z_n)^2 | \mathcal{G}_n \right\} \\ &= \frac{(n+1)^2}{S_n^2} E \{ B_{n+1}^2 (X_{n+1} - Z_n)^2 | \mathcal{G}_n \} = \frac{(n+1)^2}{S_n^2} E(B_{n+1}^2) E \{ (X_{n+1} - Z_n)^2 | \mathcal{G}_n \} \\ &= \frac{(n+1)^2}{S_n^2} E(B_{n+1}^2) Z_n (1 - Z_n) \xrightarrow{a.s.} \frac{qZ(1-Z)}{m^2} = \sigma^2. \end{aligned}$$

Since the events A_n are eventually true, one similarly obtains

$$\begin{aligned} E(V_{n+1} | \mathcal{G}_n) &\geq I_{A_n} (n+1)^2 E \left\{ \frac{B_{n+1}^2}{(S_n + c)^2} (X_{n+1} - Z_n)^2 | \mathcal{G}_n \right\} \\ &= I_{A_n} \frac{(n+1)^2}{(S_n + c)^2} E \{ B_{n+1}^2 (X_{n+1} - Z_n)^2 | \mathcal{G}_n \} \xrightarrow{a.s.} \sigma^2. \end{aligned}$$

Hence, $E(V_{n+1} | \mathcal{G}_n) \xrightarrow{a.s.} \sigma^2$. This proves condition (4) and concludes the proof of the lemma. $\square$

Theorem 2 is a quick consequence of Lemma 10. Define in fact

$$D_n = \sqrt{n}(Z_n - Z) \quad \text{and} \quad F_n = \prod_{i=n}^{\infty} (1 + H_i).$$

Because of Lemma 10,

$$\sqrt{n}(F_n Z_n - Z) = T \sqrt{n}(W_n - W) \longrightarrow \mathcal{N}(0, \sigma^2)$$

conditionally a.s. with respect to $\mathcal{G}$.

If $L < Z < U$ a.s., then $L < Z_n < U$ eventually, which in turn implies $F_n = 1$ and $D_n = \sqrt{n}(F_n Z_n - Z)$ eventually. Thus $D_n \longrightarrow \mathcal{N}(0, \sigma^2)$, conditionally a.s. with respect to $\mathcal{G}$, provided $L < Z < U$ a.s.

Lemma 11. $P(L < Z < U) = 1$.

Proof. We just prove $P(Z = L) = 0$. The proof of $P(Z = U) = 0$ is the same. Since $P(Z = L = 0) \leq P(Z = 0) = 0$, it suffices to show that $P(Z = L > 0) = 0$. Let $H = \{Z = L > 0\}$. Toward a contradiction, suppose $P(H) > 0$ and define $Q(\cdot) = P(\cdot | H)$. Since $\sqrt{n}(Z_n - L)$ is $\mathcal{G}_n$ -measurable and

$$D_n = \sqrt{n}(Z_n - Z) \leq \sqrt{n}(Z_n F_n - Z) \quad \text{if } F_n \geq 1,$$

then

$$\begin{aligned} I_{\{\sqrt{n}(Z_n - L) \leq 0\}} &= Q(D_n \leq 0 | \mathcal{G}_n) \geq Q(F_n \geq 1, \sqrt{n}(Z_n F_n - Z) \leq 0 | \mathcal{G}_n) \\ &\geq Q(\sqrt{n}(Z_n F_n - Z) \leq 0 | \mathcal{G}_n) - Q(F_n < 1 | \mathcal{G}_n) \quad \text{a.s.} \end{aligned}$$

Since $Z_n < U$ eventually on H , then $F_n \geq 1$ eventually on H . Hence, the martingale convergence theorem in [6] yields $Q(F_n < 1 | \mathcal{G}_n) \xrightarrow{Q\text{-a.s.}} 0$. By Lemma 6 and $\sigma^2 > 0$ a.s., it follows that

$$Q(\sqrt{n}(Z_n F_n - Z) \leq 0 | \mathcal{G}_n) \xrightarrow{Q\text{-a.s.}} \mathcal{N}(0, \sigma^2)((-\infty, 0]) = 1/2.$$

Thus, $I_{\{\sqrt{n}(Z_n - L) \leq 0\}} \xrightarrow{Q\text{-a.s.}} 1$, namely, $Z_n \leq L$ eventually on H , which implies the contradiction $Z_n \xrightarrow{\text{a.s.}} 1$ on H . Thus, $P(H) = 0$. $\square$

It remains only to show that Z has non-atomic distribution. This follows from the same argument of Lemma 11. Suppose in fact $P(Z = z) > 0$ for some $z \in (0, 1)$ and define $Q(\cdot) = P(\cdot | Z = z)$. Then, on the complement of a Q -null set, one obtains the contradiction $\sigma^2 = qz(1 - z)/m^2 > 0$ and

$$\delta_{\sqrt{n}(Z_n - z)}(\cdot) = Q(D_n \in \cdot | \mathcal{G}_n) \xrightarrow{\text{weakly}} \mathcal{N}(0, \sigma^2).$$

This concludes the proof of Theorem 2.

4. CONCLUDING REMARKS

Even if conceptually simple, our proofs of Theorems 1-2 are quite long and technical. This is the main reason for assuming $R_n = B_n$. However, in case $L = 0$ and $U = 1$, such assumption may be weakened into

$$E(R_n) = E(B_n) \quad \text{for all } n \geq 1;$$

see [5, Corollary 3]. Also, up to minor complications, various points in the proofs of Theorems 1-2 seem to run under the only assumption that $E(R_n) = E(B_n)$. Thus, we conjecture that Theorems 1-2 are still valid if $R_n = B_n$ is replaced by $E(R_n) = E(B_n)$.

Let

$$\overline{X}_n = (1/n) \sum_{i=1}^n X_i \quad \text{and} \quad C_n = \sqrt{n}(\overline{X}_n - Z_n).$$

If $L = 0$ and $U = 1$, as shown in [5, Corollary 3], one obtains

$$C_n \longrightarrow \mathcal{N}(0, \sigma^2 - Z(1 - Z)) \quad \text{stably.}$$

Thus, one could investigate the asymptotic behavior of C_n , or even of the pair (C_n, D_n) , in the general case $0 \leq L < U \leq 1$. Again, this could be (tentatively) performed assuming $E(R_n) = E(B_n)$ instead of $R_n = B_n$.

Another (obvious) improvement is considering multicolor urns rather than 2-colors urns. Indeed, most additional problems arising in the multicolor case are of the notational type.

A last (but minor) development could be generalizing the framework in Subsection 1.1. For instance, $B_n \vee R_n \leq c$ could be replaced by a suitable moment condition. Or else, (B_n, R_n) independent of $\sigma(\mathcal{G}_{n-1}, X_n)$ could be replaced by (B_n, R_n) conditionally independent of X_n given $\mathcal{G}_{n-1}$.

REFERENCES

- [1] Aletti G., May C., Secchi P. (2009) A central limit theorem, and related results, for a two-color randomly reinforced urn, *Adv. Appl. Probab.*, 41, 829-844.
- [2] Aletti G., Ghiglietti A., Paganoni A.M. (2013) Randomly reinforced urn designs with pre-specified allocations, *J. Appl. Probab.*, 50, 486-498.
- [3] Bai Z.D., Hu F. (2005) Asymptotics in randomized urn models, *Ann. Appl. Probab.*, 15, 914-940.
- [4] Berti P., Crimaldi I., Pratelli L., Rigo P. (2010) Central limit theorems for multicolor urns with dominated colors, *Stoch. Proc. Appl.*, 120, 1473-1491.
- [5] Berti P., Crimaldi I., Pratelli L., Rigo P. (2011) A central limit theorem and its applications to multicolor randomly reinforced urns, *J. Appl. Probab.*, 48, 527-546.
- [6] Blackwell D., Dubins L.E. (1962) Merging of opinions with increasing information, *Ann. Math. Statist.*, 33, 882-886.
- [7] Chauvin B., Pouyanne N., Sahnoun R. (2011) Limit distributions for large Polya urns, *Ann. Appl. Probab.*, 21, 1-32.
- [8] Crimaldi I., Letta G., Pratelli L. (2007) A strong form of stable convergence, *Sem. de Probab. XL, LNM*, 1899, 203-225.
- [9] Crimaldi I. (2009) An almost sure conditional convergence result and an application to a generalized Polya urn, *Internat. Math. Forum*, 4, 1139-1156.
- [10] Laruelle S., Pages G. (2013) Randomized urn models revisited using stochastic approximation, *Ann. Appl. Probab.*, 23, 1409-1436.
- [11] Mahmoud H. (2008) *Polya urn models*, Chapman-Hall, Boca Raton, Florida.
- [12] May C., Flournoy N. (2009) Asymptotics in response-adaptive designs generated by a two-color, randomly reinforced urn, *Ann. Statist.*, 37, 1058-1078.
- [13] Zhang L. (2014) A Gaussian process approximation for two-color randomly reinforced urns, *Electron. J. Probab.*, 19, 1-19.
- [14] Zhang L., Hu F., Cheung S.H., Chan W.S. (2014) Asymptotic properties of multicolor randomly reinforced Polya urns, *Adv. Appl. Probab.*, 46, 585-602.

PATRIZIA BERTI, DIPARTIMENTO DI SCIENZE FISICHE, INFORMATICHE E MATEMATICHE, UNIVERSITA' DI MODENA E REGGIO-EMILIA, VIA CAMPI 213/B, 41100 MODENA, ITALY
E-mail address: patrizia.berti@unimore.it

IRENE CRIMALDI, IMT INSTITUTE FOR ADVANCED STUDIES, PIAZZA SAN PONZIANO 6, 55100 LUCCA, ITALY
E-mail address: irene.crimaldi@imtlucca.it

LUCA PRATELLI, ACCADEMIA NAVALE, VIALE ITALIA 72, 57100 LIVORNO, ITALY
E-mail address: pratel@mail.dm.unipi.it

PIETRO RIGO (CORRESPONDING AUTHOR), DIPARTIMENTO DI MATEMATICA "F. CASORATI", UNIVERSITA' DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY
E-mail address: pietro.rigo@unipv.it